

DIOXIN DERIVATIVES AND METHOD OF MEASUREMENT THEREWITH

Patent number: EP1256579

Publication date: 2002-11-13

Inventor: KODAIRA TSUKASA (JP); YANAIHARA NOBORU (JP)

Applicant: OTSUKA PHARMA CO LTD (JP)

Classification:

- international: C07D319/24; G01N33/53

- european: C07D319/24; G01N33/53F; G01N33/531

Application number: EP20010904503 20010216

Priority number(s): WO2001JP01118 20010216; JP20000039914 20000217

Also published as:



WO0160812 (A1)

US2003108955 (A1)

CA2400840 (A1)


Abstract of EP1256579

The invention provides a highly sensitive measurement and detection technique for dioxins, which has advantages inherent to immunoassay; and markers to be used in the technique.

Disclosed are a biotinylated dioxin derivative of formula (1):

<CHEM>

(wherein X represents a hydrogen atom or a chlorine atom; R<1> represents a biotin residue; R<2>'s, which may be identical to or different from one another, independently and individually represent an arginine residue or a lysine residue; n is an integer from 1 to 5 inclusive; and m is an integer from 1 to 3 inclusive), and an immunoassay method for dioxins characterized by using the derivatives as a marker.



Data supplied from the esp@cenet database - Worldwide

DIOXIN DERIVATIVES AND METHOD OF MEASUREMENT THEREWITHClaims of **EP1256579**

1. A biotinylated dioxin derivative of formula (1):

EMI23.1

wherein X represents a hydrogen atom or a chlorine atom; R<1> represents a biotin residue; R<2>'s, which may be identical to or different from one another, independently and individually represent an arginine residue or a lysine residue; n is an integer from 1 to 5 inclusive; and m is an integer from 1 to 3 inclusive.

2. The biotinylated dioxin derivative according to claim 1, wherein X represents a hydrogen atom, n is 4, and M is 2.

3. An immunoassay method for dioxins characterized by comprising using as a marker a biotinylated dioxin derivative of formula (1):

EMI23.2

wherein X represents a hydrogen atom or a chlorine atom; R<1> represents a biotin residue; R<2>'s, which may be identical to or different from one another, independently and individually represent an arginine residue or a lysine residue; n is an integer from 1 to 5 inclusive; and m is an integer from 1 to 3 inclusive.

4. A reagent for immunoassay of dioxins, containing a biotinylated dioxin derivative of formula (1):

EMI24.1

wherein X represents a hydrogen atom or a chlorine atom; R<1> represents a biotin residue; R<2>'s, which may be identical to or different from one another, independently and individually represent an arginine residue or a lysine residue; n is an integer from 1 to 5 inclusive; and m is an integer from 1 to 3 inclusive.

Data supplied from the **esp@cenet** database - Worldwide